

Lecture 4: Sea-floor depth, age, and heat flow


- How do we map the seafloor today?
 - Gravity, strain, and the geoid
- Stochastic reheating model



We acknowledge and respect the lək̓ʷəŋən peoples on whose traditional territory the university stands and the Songhees, Esquimalt and W̱SÁNEĆ peoples whose historical relationships with the land continue to this day.




Mapping the sea-floor



How do we map the bathymetry of the sea-floor?



Mapping the sea-floor




How do we map the bathymetry of the sea-floor?

A combination of:


- Depth Soundings
- Satellite Altimetry



Bathymetric Prediction From SEASAT Altimeter Data (Dixon et al. 1983)



Bathymetric Prediction From SEASAT Altimeter Data (Dixon et al. 1983)



Gravitational potential

Gravitational potential is the the work (energy transferred) per unit mass that would be needed to move an object to that point from a distance infinitely far away. Recall that: $\text{work} = \text{force} \times \text{displacement}$



Gravitational potential

Gravitational potential is the the work (energy transferred) per unit mass that would be needed to move an object to that point from a distance infinitely far away. Recall that: $\text{work} = \text{force} \times \text{displacement}$


The acceleration due to gravity is constant along an **equipotential surface**. [How is the geoid related to gravitational potential?](#)



Gravitational potential

Gravitational potential is the the work (energy transferred) per unit mass that would be needed to move an object to that point from a distance infinitely far away. Recall that: $\text{work} = \text{force} \times \text{displacement}$


The acceleration due to gravity is constant along an **equipotential surface**. [How is the geoid related to gravitational potential?](#)



Gravitational potential

Gravitational potential is the the work (energy transferred) per unit mass that would be needed to move an object to that point from a distance infinitely far away. Recall that: $\text{work} = \text{force} \times \text{displacement}$

The acceleration due to gravity is constant along an **equipotential surface**. [How is the geoid related to gravitational potential?](#)



The **geoid** (a model) is the ocean surface elevation if winds and tides were absent.



If the Earth is isostatically compensated, why does the geoid (sea-surface) vary?



If the Earth is isostatically compensated, why does the geoid (sea-surface) vary?

Strain in the asthenosphere removes any stress that exists due to force imbalances (ie density contrasts).. so geoid variations must:

- be small enough that the lithosphere can hold that stress
- be caused by other dynamic forces:
 - convection/dynamic topography
 - topography that changes faster than the asthenosphere can compensate (faster than the strain rate)



If the Earth is isostatically compensated, why does the geoid (sea-surface) vary?




If the Earth is isostatically compensated, why does the geoid (sea-surface) vary?

Consider (sketch and calculate) the acceleration due to gravity above Hawaii's big island (*for this back-of-the-envelope calculation, you can assume the masses are point-masses*). Gravitational potential energy $\propto$ force due to gravity and

$$F = -G \frac{m_1 m_2}{r^2} \quad F = m \times a \quad a = -G \frac{m_1}{r^2}$$

- $G = 6.6743 \times 10^{-11} \frac{\text{m}^3}{\text{kg s}^2}$
- Mass of the big island: $1.6 \times 10^{15} \text{ kg}$
- Mass of Earth: $5.97 \times 10^{24} \text{ kg}$
- Radius of Earth: $6.378 \times 10^6 \text{ m}$
- Depth of seafloor: $6 \times 10^3 \text{ m}$

Assume that the island is exactly 6 km tall. How is the acceleration due to gravity at sea level different above this island when compared to some far away location in the Pacific that has no island? Is it higher/lower? How much so?




If the Earth is isostatically compensated, why does the geoid (sea-surface) vary?

Consider (sketch and calculate) the acceleration due to gravity above Hawaii's big island (*for this back-of-the-envelope calculation, you can assume the masses are point-masses*). Gravitational potential energy $\propto$ force due to gravity and

$$F = -G \frac{m_1 m_2}{r^2} \quad F = m \times a \quad a = -G \frac{m_1}{r^2}$$

- $G = 6.6743 \times 10^{-11} \frac{\text{m}^3}{\text{kg s}^2}$
- Mass of the big island: $1.6 \times 10^{15} \text{ kg}$
- Mass of Earth: $5.97 \times 10^{24} \text{ kg}$
- Radius of Earth: $6.378 \times 10^6 \text{ m}$
- Depth of seafloor: $6 \times 10^3 \text{ m}$

Assume that the island is exactly 6 km tall. How is the acceleration due to gravity at sea level different above this island when compared to some far away location in the Pacific that has no island? Is it higher/lower? How much so? **Now consider the force 3 km away from the island. What percent of the force is felt at this distance?**



If the Earth is isostatically compensated, why does the geoid (sea-surface) vary?

Consider (sketch and calculate) the acceleration due to gravity above Hawaii's big island (*for this back-of-the-envelope calculation, you can assume the masses are point-masses*). Gravitational potential energy $\propto$ force due to gravity and

$$F = -G \frac{m_1 m_2}{r^2} \quad F = m \times a \quad a = -G \frac{m_1}{r^2}$$

- $G = 6.6743 \times 10^{-11} \frac{\text{m}^3}{\text{kg s}^2}$
- Mass of the big island: $1.6 \times 10^{15} \text{ kg}$
- Mass of Earth: $5.97 \times 10^{24} \text{ kg}$
- Radius of Earth: $6.378 \times 10^6 \text{ m}$
- Depth of seafloor: $6 \times 10^3 \text{ m}$

Assume that the island is exactly 6 km tall. How is the acceleration due to gravity at sea level different above this island when compared to some far away location in the Pacific that has no island? Is it higher/lower? How much so? **Now consider the force 3 km away from the island. What percent of the force is felt at this distance? Repeat this same exercise using a point mass that is on the sea-floor (ie deeper than the previous calculation). The exact mass here doesn't matter. What percent of the force due to that mass is experienced 3 km away on the sea surface?**

In [11]:

```
1 import numpy as np
2 from matplotlib import pyplot as plt
3 G=6.6743e-11# m3 kg-1 s-2
4 ## point mass at 0,-6km
```

If the Earth is isostatically compensated, why does the geoid (sea-surface) vary? (Summary)




If the Earth is isostatically compensated, why does the geoid (sea-surface) vary? (Summary)

- small amplitude differences in gravitational potential are maintained by lithosphere strength
- due to lateral differences in density (ie ocean at 3km next to basalt at 3km)
- low frequency (long wavelength) variations in gravitational potential are due to difference deeper in the Earth
- high frequency (short wavelength) variations are due to differences near the surface of the Earth




Mapping the sea-floor



- Altimetry data decomposed into high frequency (low wavelength) and low frequency (high wavelength) *spectral* components
- Low frequency components are combined with ship soundings to estimate deeper Earth density structures
- High frequency components are used to interpolate between soundings, assuming deep structure constant
- High frequency components are used to resolve very shallow density contrasts (such as sea-mounts)



Mapping the sea-floor



- Altimetry data decomposed into high frequency (low wavelength) and low frequency (high wavelength) *spectral* components
- Low frequency components are combined with ship soundings to estimate deeper Earth density structures
- High frequency components are used to interpolate between soundings, assuming deep structure constant
- High frequency components are used to resolve very shallow density contrasts (such as sea-mounts)



Returning to the plate model and old oceanic lithosphere



(left) A and B show sea-floor depth (km) versus age (0 to 180 Ma). Lines are three different "plate" models. (right) Histogram of depth. Plate models are the dashed and dotted lines. "Random reheating" model is the solid line.



Returning to the plate model and old oceanic lithosphere

