


Contents ↻ ⚙

- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
 - ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
 - Constant crustal thickness
 - Constant crustal thickness

Lecture 5: Hot spots and seamounts

The topography and geology of the seafloor offers some of the only clues we have to understand the hidden workings of the mantle below. We have discussed some of the largest features of ocean today we consider the small seamounts and their critical role in this story.

- Models of mantle convection and implications for volcanic islands
- Plumes and hotspots
- Free air anomalies and the **swell**
 - Explaining swells (thick crust vs thin lithospheric mantle)



We acknowledge and respect the lək̓ʷəŋən peoples on whose traditional territory the university stands and the Songhees, Esquimalt and W̱S̱X̱ historical relationships with the land continue to this day.

Features of the seafloor

Contents ↻ ⚙

▼ Lecture 5: Hot spots and seamounts

Features of the seafloor

▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)

Possible convective models

The volcanic consequences of each convective model

The volcanic consequences of each convective model

The source of lavas must be deep

The plume

The ages of "rising bubbles"

More gravity..

More gravity..

More gravity..

More gravity..

More gravity..

Swells and depth anomalies


▼ Swells and depth anomalies

How are swells supported?

How are swells supported?

Constant crustal thickness

Constant crustal thickness




Contents ↻ ⚙

- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
 - ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
 - Constant crustal thickness
 - Constant crustal thickness

A possible origin of the Hawaiiaian Islands (Tuzo Wilson, 1963)

LEGEND

- Median Ridge
- Flow Direction
- Island Chain
- - - Fault



*"It is less obvious why a stream of volca
arise like a series of bubbles from a poin
island of Hawaii which is far from the ri*
**Tuzo Wilson, A possible origin of the H
1963**

Contents ↻ ⚙

▼ Lecture 5: Hot spots and seamounts

Features of the seafloor

▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)

Possible convective models

The volcanic consequences of each convective model

The volcanic consequences of each convective model

The source of lavas must be deep

The plume

The ages of "rising bubbles"

More gravity..

More gravity..

More gravity..

More gravity..

More gravity..

Swells and depth anomalies

▼ Swells and depth anomalies

How are swells supported?

How are swells supported?

Constant crustal thickness

Constant crustal thickness

Possible convective models




FIG. 2. Three possible modes of convection in the Earth's

Contents ↻ ⚙

- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
 - ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
 - Constant crustal thickness
 - Constant crustal thickness

The volcanic consequences of each convective model




FIG. 3. Some possible patterns of convection, showing that, if active volcanoes are formed by rising vertical currents, chains of extinct volcanoes might be formed by the horizontal movement of the currents. The shaded areas represent stable cores of cells.

Contents ↻ ⚙

- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
 - ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
 - Constant crustal thickness
 - Constant crustal thickness

The volcanic consequences of each convective model



FIG. 4. Diagram to illustrate that in general if lava is generated at some other convecting system than over a rising current, then only one volcano will be generated.

The source of lavas must be deep

Contents ↻ ⚙

▼ Lecture 5: Hot spots and seamounts

Features of the seafloor

▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)

Possible convective models

The volcanic consequences of each convective model

The volcanic consequences of each convective model

The source of lavas must be deep

The plume

The ages of "rising bubbles"

More gravity..

More gravity..

More gravity..

More gravity..

More gravity..

Swells and depth anomalies

▼ Swells and depth anomalies

How are swells supported?

How are swells supported?

Constant crustal thickness

Constant crustal thickness




FIG. 5. Diagram to illustrate that if lava is generated in the stable core of a convection cell and the surface is carried by the jet stream, then one source can give rise to a chain of volcanoes even if the source is not over a rising current. This is proposed as a possible origin of the Hawaiian chain of islands.

Contents ↻ ⚙

- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
- ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
- Constant crustal thickness
- Constant crustal thickness

The plume


"In my model there are about twenty deep mantle plumes bringing heat and relatively primordial material to the surface. The convection cells in the asthenosphere flow radially away from each of these plumes. This model is compatible with the observation that there is a difference between oceanic island arc basalts and continental basalts."

Jason Morgan, Convection Plumes in the Lower Mantle

The ages of "rising bubbles"

Contents ↻ ⚙

- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
 - ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
 - Constant crustal thickness
 - Constant crustal thickness



K-Ar ages from Hawaii chain volcanoes (Dalrymple et. al. 1977)

Contents ↻ ⚙

- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
- ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
- Constant crustal thickness
- Constant crustal thickness



More gravity..

Gravity measurements are small variations in the acceleration due to gravity (similar to the w you did from last lecture). [How do we actually make these measurements?](#)

More gravity..

Contents ↻ ⚙

▼ Lecture 5: Hot spots and seamounts

Features of the seafloor

▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)

Possible convective models

The volcanic consequences of each convective model

The volcanic consequences of each convective model

The source of lavas must be deep

The plume

The ages of "rising bubbles"

More gravity..

More gravity..

More gravity..

More gravity..

More gravity..

Swells and depth anomalies

▼ Swells and depth anomalies


How are swells supported?

How are swells supported?

Constant crustal thickness

Constant crustal thickness

Gravity measurements are small variations in the acceleration due to gravity (similar to the w you did from last lecture). [How do we actually make these measurements?](#)



Pendulum of Sterneck 1887

The equation of simple harmonic motion, $T = 2\pi\sqrt{\frac{L}{g}}$, describes the relation between an easily measured property (T, the period of oscillation) to *gravity*, g.

Energy is added to the pendulum in a regular fashion to compensate for friction.

More gravity..

Contents ↻ ⚙

▼ Lecture 5: Hot spots and seamounts

Features of the seafloor

▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)

Possible convective models

The volcanic consequences of each convective model

The volcanic consequences of each convective model

The source of lavas must be deep

The plume

The ages of "rising bubbles"

More gravity..

More gravity..

More gravity..

More gravity..

More gravity..

Swells and depth anomalies

▼ Swells and depth anomalies


How are swells supported?

How are swells supported?

Constant crustal thickness

Constant crustal thickness

Gravity measurements are small variations in the acceleration due to gravity (similar to the way you did from last lecture). [How do we actually make these measurements?](#)



Pendulum of Sterneck 1887

The equation of simple harmonic motion, $T = 2\pi\sqrt{\frac{L}{g}}$, describes the relation between an easily measured property (T, the period of oscillation) to *gravity*, g.


Energy is added to the pendulum in a regular fashion to compensate for friction.

[We keep talking about gravity measurements in the ocean.. how were these measurements made on a ship?](#)

Contents ↻ ⚙

- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
- ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
- Constant crustal thickness
- Constant crustal thickness

More gravity..



Shipboard gravimeter built in 1923

The equation of simple harmonic motion

$T = 2\pi\sqrt{\frac{L}{g}}$, describes the relation

an easily measured property (T, the period of oscillation) to *gravity, g*.

Shipboard accelerations are certainly not simple harmonic motion, however...

By precisely measuring the deflection of multiple pairs of pendulums, gravity can be compared under additional accelerations. A set of multiple pendulums are needed to account for

Contents ↻ ⚙

- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
- ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
- Constant crustal thickness
- Constant crustal thickness

More gravity..




Illustration of the twin GRACE satellites

Today, we mostly measure gravity using pendulums. The GRACE (Gravity Recovery and Climate Experiment) mission launched in 2002 and other missions using a similar approach are planned for the future. These satellites measure gravity up to 1000 times more accurately than pendulums and have been really important in observing gravity *change* on the Earth due to groundwater loss or ice sheet decay.

More gravity..

Contents ↻ ⚙

▼ Lecture 5: Hot spots and seamounts

Features of the seafloor

▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)

Possible convective models

The volcanic consequences of each convective model

The volcanic consequences of each convective model

The source of lavas must be deep

The plume

The ages of "rising bubbles"

More gravity..

More gravity..

More gravity..

More gravity..

More gravity..

Swells and depth anomalies


▼ Swells and depth anomalies

How are swells supported?

How are swells supported?

Constant crustal thickness

Constant crustal thickness



Free air gravity anomaly around Hawaii (~1000 km; Figure from Dunn et. al. 2024). Recall gravity anomalies are only corrected for the elevation difference from sea-level of the measurement. In practice, a specific geoid is used and it may not be perfect, but this **anomaly** highlights gravity due to more (positive anomaly) or less (negative anomaly) mass than expected beneath the

Swells and depth anomalies

Contents ↻ ⚙

▼ Lecture 5: Hot spots and seamounts

Features of the seafloor

▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)

Possible convective models

The volcanic consequences of each convective model

The volcanic consequences of each convective model

The source of lavas must be deep

The plume

The ages of "rising bubbles"

More gravity..

More gravity..

More gravity..

More gravity..

More gravity..

Swells and depth anomalies


▼ Swells and depth anomalies

How are swells supported?

How are swells supported?

Constant crustal thickness

Constant crustal thickness




Bathymetry of the Pacific. Transect #5 is the next figure. (Watts 1976).

Swells and depth anomalies

Contents ↻ ⚙

- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
- ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
 - Constant crustal thickness
 - Constant crustal thickness




Transects across Hawaii (Crough 1983)

How are swells supported?

Contents ↻ ⚙


- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
- ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
- Constant crustal thickness
- Constant crustal thickness



How are swells supported?

Contents ↻ ⚙

- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
- ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
 - Constant crustal thickness
 - Constant crustal thickness



Constant crustal thickness

Contents ↻ ⚙

▼ Lecture 5: Hot spots and seamounts

Features of the seafloor

▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)

Possible convective models

The volcanic consequences of each convective model

The volcanic consequences of each convective model

The source of lavas must be deep

The plume

The ages of "rising bubbles"

More gravity..

More gravity..

More gravity..

More gravity..

More gravity..

Swells and depth anomalies


▼ Swells and depth anomalies

How are swells supported?

How are swells supported?

Constant crustal thickness

Constant crustal thickness




Dunn et. al. 2024

Constant crustal thickness

Contents ↻ ⚙

- ▼ Lecture 5: Hot spots and seamounts
 - Features of the seafloor
 - ▼ A possible origin of the Hawaiian Islands (Tuzo Wilson, 1963)
 - Possible convective models
 - The volcanic consequences of each convective model
 - The volcanic consequences of each convective model
 - The source of lavas must be deep
 - The plume
 - The ages of "rising bubbles"
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - More gravity..
 - Swells and depth anomalies
- ▼ Swells and depth anomalies
 - How are swells supported?
 - How are swells supported?
- Constant crustal thickness
- Constant crustal thickness



Dunn et. al. 2024