


# Lecture 7: Hot spots, seamounts, and ridges

The topography and geology of the seafloor offers some of the only clues we have to understanding the hidden workings of the mantle below. We have discussed some of the largest features of ocean basins, and today we consider the small seamounts and their critical role in this story.

- Axial depth and crustal thickness
  - $\text{Na}_2\text{O}$  in MORB
- Spreading rate vs potential temperatures
- Trace elements and isotopes in MORB/OIB



*We acknowledge and respect the  $\text{lək}^w\text{əŋən}$  peoples on whose traditional territory the university stands and the Songhees, Esquimalt and  $\text{W̱SÁNEĆ}$  peoples whose historical relationships with the land continue to this day.*




## *Review*

What evidence do we have for mantle convection?

- Magnetic anomalies that are symmetric around MOR
- Elevation of seafloor fits diffusive cooling model (up to ~50 Ma)
- Constant crustal thickness in the oceanic crust
- Higher heatflow under MOR than far from MOR
- Ocean basins are young

What evidence do we have for plumes?

- Anomalously high elevation swell surrounding active mid-plate (usually) volcanism
- Often a monotonic age progression of volcanism the youngs in the plate-motion direction



## *Review*

What evidence do we have for mantle convection?

- Magnetic anomalies that are symmetric around MOR
- Elevation of seafloor fits diffusive cooling model (up to ~50 Ma)
- Constant crustal thickness in the oceanic crust
- Higher heatflow under MOR than far from MOR
- Ocean basins are young

What evidence do we have for plumes?

- Anomalously high elevation swell surrounding active mid-plate (usually) volcanism
- Often a monotonic age progression of volcanism the youngs in the plate-motion direction

Today we will go into some of the geochemical properties of mid-ocean ridge and ocean island basalts to get a better understanding of how the mantle on Earth works.



## Axial depth, crustal thickness, and melting





FIG 7 (a) Adiabatic decompression paths calculated using the equations given by McKenzie (1984a) Appendix D, a fourth order Runge-Kutta scheme and

$$\Delta S = 250 \text{ J kg}^{-1} \text{ } ^\circ\text{C}^{-1}.$$

The curves are labelled with their **potential** temperatures, and entropy is conserved to 1 part in  $10^4$  during the numerical integration. The curves between the solidus and the liquidus are labelled with the melt fraction by weight. (b) The total thickness of melt present below a given depth plotted as a function of depth, calculated by integrating the volume of melt present in (a).




*Melting peridotite*





## *Peridotite melting experiments*

(-0.5 is the solidus, 0.5 is the liquidus)




## *Peridotite melting experiments*

(-0.5 is the solidus, 0.5 is the liquidus)




## *Na<sub>2</sub>O data from MORB*



What is controlling the Na to Mg ratio?





## Na<sub>2</sub>O data from MORB



What is controlling the Na to Mg ratio? Percent melting AND differentiation. Na<sub>2</sub>O is controlled by percent partial melting but the magmas can lose phases during ascent. Olivines or pyroxenes may literally *drop out* of the rising magma, changing the major element chemistry. By *normalizing* Na<sub>2</sub>O data to a fixed MgO content, percent melting (potential temperature) can be isolated.



Normalized (MgO = 8%) Na<sub>2</sub>O




## Variations in spreading rate?

If melting is *passive*, what is the expected relationship between spreading rate and axial depth?




## Variations in spreading rate?

If melting is *passive*, what is the expected relationship between spreading rate and axial depth?



## *Potential temperatures beneath ridges and oceanic islands*

The partitioning of Mg in Olivine is sensitive to temperature while the Fe/Mg partitioning is not




Strong geochemical evidence for the existence of thermally driven mantle plumes



Trace elements in MORB and OIB (review compatibility trends)



## Trace elements in MORB and OIB (review compatibility trends)



Tristan, South Atlantic




Tubuai, French Polynesia


Trace elements in MORB and OIB (review compatibility trends)



Trace elements in MORB and OIB (review compatibility trends)



# Models of the mantle



## Sm and Rb

- $^{147}\text{Sm} \rightarrow ^{143}\text{Nd}$ 
  - Sm more **compatible** than Nd
- $^{87}\text{Rb} \rightarrow ^{87}\text{Sr}$ 
  - Rb highly **incompatible** (and more incompatible than Sr)
- What trends do you expect between primitive mantle, MORB, and continental crust?
  - If plumes sample primitive mantle, what should their radiogenic Nd and Sr look like?
  - *sketch:*  $\frac{^{87}\text{Sr}}{^{86}\text{Sr}}$  (x) vs  $\frac{^{143}\text{Nd}}{^{144}\text{Nd}}$  (y)

