





Lecture 8: Stratigraphic time part I

1. Gross vs. net fluxes
 - A. erosion and hiatus
 - B. how to build a rock record
2. Correlative surfaces
 - A. Chronostratigraphy (Wheeler diagrams)
3. Time in the rock record
 - A. Sadler effect


We acknowledge and respect the $l\acute{a}k^{w}\acute{\eta}n$ peoples on whose traditional territory the university stands and the Songhees, Esquimalt and $W\acute{S}\acute{A}N\acute{E}\acute{C}$ peoples whose historical relationships with the land continue to this day.




Gross and net sedimentation




Gross and net sedimentation




Gross and net sedimentation




Gross and net sedimentation




Gross and net sedimentation




Gross and net sedimentation



Gross and net sedimentation




Gross and net sedimentation



net sedimentary record is full of **gaps**: both **erosive surfaces** and **hiatuses**.



Gross and net sedimentation



net sedimentary record is full of **gaps**: both **erosive surfaces** and **hiatuses**.

How much time is represented by sedimentary rock?



How to build the net rock record

```
In [2]: import csv
import numpy as np
from matplotlib import pyplot as plt
import seaborn as sns
sns.set_context('talk')
%matplotlib inline

#empty lists for data
model_topo=[]

#open the data file
▼ with open("model_results.csv", "r") as fid:
    data = csv.reader(fid, delimiter=",")

    #reads the data in line-by-line
    ▼ for row in data:
        model_topo.append([float(r) for r in row[1:]])


#convert to list of numpy arrays
model_topo=[np.array(t) for t in model_topo]

#define as change from initial conditions
model_topo=[t-model_topo[0] for t in model_topo]
```




How to build the net rock record

```
In [3]: #pull topographic evolution from one spot in the basin  
section=[t[454] for t in model_topo]  
  
plt.plot(section, '-', color='r')  
plt.plot(np.arange(len(section))[0::4], section[0::4], '.', markersize=5, color='k')  
  
plt.gca().set_xticks([])  
plt.gca().set_xlabel(r'time $\rightarrow$')  
plt.gca().set_ylabel('height (meters)')  
plt.gca().xaxis.set_label_position('top')
```




How to build the net rock record

```
In [3]: net=np.flipud(section) #flip this stratigraphy upside down to examine youngest to oldest
▼ for i,s in enumerate(net): #count through time slices starting at the youngest
▼   if i!=len(net)-1: #if we're not at the end
      #look "down section" - was this layer eroded?
      # --> if height had decreased, then yes
▼   if s<net[i+1]: #s is height of current timeslice, net[i+1] is heigh of previous timeslice
      net[i+1]=s #create new record of net sedimentation
#make stratigraphy younging upwards again
net=np.flipud(net)
plt.plot(net,color='b')
plt.fill_between(np.arange(len(section)),section,net,color='r',alpha=0.25)
plt.gca().set_xticks([])
plt.gca().set_xlabel(r'time $\rightarrow$')
plt.gca().set_ylabel('height (meters)')
plt.gca().xaxis.set_label_position('top')
```




How to build the net rock record




Coastline position and erosion and hiatus

We will look at a set of model runs to better understand the relationship of the coastline position to **erosion** and **hiatus** (and **time**). First, let's review the controls on coastline position:


- Coastline progrades/regresses/moves seaward when: $\frac{sed}{dt} > \frac{space}{dt}$
- Coastline retrogrades/transgresses/moves landward when: $\frac{sed}{dt} < \frac{space}{dt}$
- Coastline aggrades/stays in the same location landward when: $\frac{sed}{dt} \approx \frac{space}{dt}$




Simple sea level cycles: stratigraphic columns



Simple sea level cycles: time vs topography




Simple sea level cycles: time vs topography




Where do you get erosion? Where is erosion rate maximized? What about non-deposition? (*annotate/draw on your handout*)




Simple sea level cycles: time vs topography (erosion)




Simple sea level cycles: time vs topography (**erosion** and hiatus)



Simple sea level cycles: *Wheeler* diagram




Simple sea level cycles: *Wheeler* diagram




Note: not all positive deposition is significant. Discuss where you think the thickest sediments (highest sedimentation rates) are.




Simple sea level cycles: stratigraphic profile




Class example 1: stratigraphic columns




Class example 1: time vs topography




Class example 1: time vs topography (**erosion**)




Class example 1: time vs topography (**erosion** and hiatus)




Class example 1: *Wheeler* diagram




Class example 1: stratigraphic profile




Class example 2: stratigraphic columns




Class example 2: time vs topography




Class example 2: time vs topography (erosion)




Class example 2: time vs topography (**erosion** and hiatus)




Class example 2: *Wheeler* diagram




Class example 2: stratigraphic profile




On/Off sediment supply with constant subsidence: time vs topography




On/Off sediment supply with constant subsidence: time vs topography




On/Off sediment supply with constant subsidence: time vs topography (**erosion**)



On/Off sediment supply with constant subsidence: time vs topography (**erosion** and hiatus)



On/Off sediment supply with constant subsidence: *Wheeler* diagram



On/Off sediment supply with constant subsidence: stratigraphic profile

